

ERRATUM

J. Raptor Res. 36(2):157

© 2002 The Raptor Research Foundation, Inc.

FACTORS INFLUENCING LENGTH OF THE POST-FLEDGING PERIOD AND TIMING OF DISPERSAL IN BONELLI'S EAGLE (*HIERAAETUS FASCIATUS*) IN SOUTHWESTERN SPAIN

ELENA ANGULO¹

Estación Biológica de Doñana, Apdo. 1056, 41080 Sevilla, Spain and Instituto de Investigación en Recursos Cinegéticos, Apdo. 535, Ciudad Real, Spain

Figures 1 and 3 printed in the article "Factors influencing length of the post-fledging period and timing of dispersal in Bonelli's Eagle (*Hieraetus fasciatus*) in southwestern Spain," (*Journal of Raptor Research* 35[3]:228–234) were incorrect, draft versions. The correct, revised figures that correspond to this article and that should be substituted for Figure 1 (page 229) and Figure 3 (page 231) are printed below.

¹ E-mail address: angulo@ebd.csic.es




Figure 1. Study area in southwestern Spain, 1998. Grey circles represent the ten territories of Bonelli's Eagles, and numbered black squares are the six sites of prey availability counts.




Figure 3. Monthly prey abundance for the six surveyed sites (see Fig. 1 for the geographic locations) during the post-fledging period. (a) European rabbit (*Oryctolagus cuniculus*) abundance. (b) Red-legged Partridge (*Alectoris rufa*) abundance.